

Fortified StimulateTM Yield Enhancer

ACTIVE INGREDIENTS:

Cytokinin (as kinetin)	0.009%
Gibberellic acid	0.005%
Indole-3-butyric acid	0.005%
Indole-3-acetic acid	0.005%

This product contains: 0.002682 grams cytokinin, 0.001490 grams gibberellic acid, 0.001490 grams indole-3-butyric acid, 0.001490 grams indole-3-acetic acid per fluid ounce.

CONTAINS NON-PLANT FOOD INGREDIENTS:

0.009% Cytokinin
0.005% Gibberellic Acid
0.005% Indole-3-butyric Acid
0.005% Indole-3-acetic Acid

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.html>

KEEP OUT OF REACH OF CHILDREN CAUTION

SEE ADDITIONAL PRECAUTIONARY STATEMENTS
AND DIRECTION FOR USE IN BOOKLET.

Z- FORTSTIMYE

Rev: 20180601

F2399

**Registration Number
2018066A Fertilizers Act**

**Manufactured and Guaranteed by
Stoller Enterprises, Inc.**

9090 Katy Freeway, Suite 400
Houston, Texas 77024 USA

Phone: 1-800-539-5283 or 713-461-1493




Stoller

Fortified Stimulate Yield Enhancer

ACTIVE INGREDIENTS:

Cytokinin (as kinetin)	0.009%
Gibberellic acid	0.005%
Indole-3-butyric acid	0.005%
Indole-3-acetic acid	0.005%

This product contains: 0.002682 grams cytokinin, 0.001490 grams gibberellic acid, 0.001490 grams indole-3-butyric acid, 0.001490 grams indole-3-acetic acid per fluid ounce.

CONTAINS NON-PLANT FOOD INGREDIENTS:

0.009% Cytokinin
0.005% Gibberellic Acid
0.005% Indole-3-butyric Acid
0.005% Indole-3-acetic Acid

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE ADDITIONAL PRECAUTIONARY STATEMENTS ON LEFT PANEL.

Registration Number 2018066A Fertilizers Act

Read attached label before using.

FIRST AID

If inhaled

- Move person to fresh air.
- If person is not breathing, call 911 or ambulance; then give artificial respiration, preferably mouth-to-mouth, if possible.
- Call a poison control center or doctor for further treatment advice.

If in eyes

- Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes; then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

If on skin or clothing

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency information concerning this product, call the National Pesticides Information Center (NPIC) at 1-800-858-7378 seven days a week, 6:30 am to 4:30 pm Pacific Time (NPIC website: www.npic.orst.edu).

Manufactured by

Stoller Enterprises, Inc.

9090 Katy Freeway, Suite 400
Houston, Texas 77024 USA

Phone: 800-539-5283
or 713-461-1493

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION

Harmful if inhaled. Avoid breathing vapor or spray mist. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Remove and wash contaminated clothing before reuse. Wear the appropriate Personal Protective Equipment (PPE).

PERSONAL PROTECTIVE EQUIPMENT

Waterproof gloves are sufficiently chemical-resistant for this product.

Applicators and other handlers must wear:

- long-sleeved shirt and long pants,
- chemical-resistance gloves made of any waterproof material such as polyethylene or polyvinyl chloride,
- shoe plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

For terrestrial uses, do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not apply where runoff is likely to occur. Do not contaminate water when cleaning equipment or disposing of equipment washwater or rinsate. Exposed treated seed may be hazardous to birds and other wildlife. Treat only those seeds needed for immediate use and planting. Do not store excess treated seed beyond planting time. Dispose of all excess treated seed and seed packaging by burial away from streams and bodies of water.

DIRECTIONS FOR USE

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural products. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval.

Do not re-enter or allow re-entry into treated areas until 12 hours after application.

PPE required for early entry to treated areas that involves contact with anything that has been treated, such as plants, soil, or water is: coveralls, chemical-resistant gloves made of any waterproof material, and shoes plus socks.

CHEMIGATION: APPLICATION AND CALIBRATION TECHNIQUES FOR SPRINKLER IRRIGATION

Apply this product only through the following types of irrigation systems: center pivot, traveler, big gun, motorized lateral move, end tow, side (wheel) roll, solid set, and hand move irrigation equipment. Do not apply through any other types of irrigation systems. Crop injury or lack of effectiveness can result from non-uniform distribution of treated water. If you have questions about calibration, you should contact Experiment Station specialists, equipment manufacturers or other experts. Do not use an irrigation system (including greenhouse systems) used for product application to a public water system unless the product label-prescribed safety devices for public water systems are in place. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person shall shut the system down and make necessary adjustments should the need arise.

A.Center Pivot, Traveler, Big Gun, Motorized Lateral Move, End Tow, and Side (Wheel) Roll Irrigation Equipment: Operate system and injection equipment at normal pressures recommended by the manufacturer of injection equipment used. Fill tank of injection equipment with water. Operate system for one complete circle for center pivot or one complete run for the other recommended equipment, measuring time required, amount of water injected, and hectareage contained in circle or run. Mix recommended amount of Fortified Stimulate Yield Enhancer for hectareage to be covered into same amount of water used during calibration and inject into system continuously for one revolution or run, but continue to operate irrigation system until Fortified Stimulate Yield Enhancer has been cleared from last sprinkler head. Spray mixture in the chemical supply tank must be agitated at all times, otherwise settling and uneven application may occur.

B.Solid Set and Hand Move Irrigation

Equipment: Determine hectareage covered by sprinkler. Fill tank of injection equipment with water and adjust flow to use contents over a thirty to forty-five minute period. Mix desired amount of Fortified Stimulate Yield Enhancer for hectareage to be covered into quantity of water used during calibration and operate entire system at normal pressures recommended by the manufacturer of injection equipment used for amount of time established during calibration. Provide constant mechanical agitation in the mix tank to insure that Fortified Stimulate Yield Enhancer will remain in suspension during the injection cycle. Fortified Stimulate Yield Enhancer can be injected at the beginning or end of the irrigation cycle or as a separate application. Stop injection equipment after treatment is completed and continue to operate irrigation system until Fortified Stimulate Yield Enhancer is cleared from last sprinkler head.

SAFETY DEVICES

(1) The systems designed above must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow. (2) All product injection pipelines must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. (3) The product injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. (4) The system must contain functional interlocking controls to automatically shut off the product injection pump when the water pump motor stops. (5) The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where product

distribution is adversely affected. (6) Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with agricultural products and capable of being fitted with a system interlock. (7) Do not apply when wind speed favors drift beyond the area intended for treatment.

SYSTEMS CONNECTED TO PUBLIC WATER SOURCES

Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of product introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to product introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe. The system must contain functional interlocking controls to automatically shut off the product injection pump when the water pump motor stops or, in cases where there is no water pump, when the water pressure decreases to the point where product distribution is adversely affected.

For additional instructions on safety precautions, refer to statements (2), (3), (4) (6) and (7) in the section SAFETY DEVICES.

GENERAL INFORMATION

Fortified Stimulate Yield Enhancer is a biostimulant containing plant growth regulators. Stimulate enhances plant growth and development by stimulating cell division, cell differentiation and enlargement, nutrient uptake and nutrient utilization. It is especially effective when applied with foliar fertilizer, but it is also compatible with agricultural products.

MIXING INSTRUCTIONS: Fortified Stimulate Yield Enhancer is water soluble and suitable for use in conventional liquid application systems including sprinkler irrigation systems. Shake Stimulate thoroughly and dilute in sufficient water to assure adequate, even coverage without producing excessive runoff. Agitate the spray mixture during application and apply within 12 hours of dilution. Stimulate can be applied tank mixed with most insecticides, fungicides, herbicides and foliar fertilizers but must be the last addition to the spray mixture.

APPLICATION INSTRUCTIONS

Apply Fortified Stimulate Yield Enhancer (by ground or air) to foliage diluted in 18.71 liters to 93.54 liters of water per hectare. Larger volumes of water may be used if not associated with excessive runoff. For best results, apply in the early morning or late evening. When applying Fortified Stimulate Yield Enhancer in a band or as a foliar-directed spray, reduce the application rate from the labeled broadcast rate in proportion to the percent of the field surface area covered by the foliar spray, but not below the minimum rate listed in the table.

VEGETABLES

Crop	No. of Applications	Broadcast Spray	Band Spray	In Seed Furrow	Transplant Water	Spray Timing
		MI/H	MI/H	MI/H	MI/H	
Beans	1	584	292			Between the 3 rd trifoliolate leaf stage and flower bud formation.
	2-3	292	219	146-584		Begin at the 3 rd trifoliolate leaf stage and then at 7-10 day intervals.
	4-6	219	146			Begin at the 2 nd trifoliolate leaf stage and then at 7-14 day intervals.
Broccoli	3	438	292	146-584	146-584	Begin at the 4-5 leaf stage and then at 10-14 day intervals.
Brussel Sprouts	3	438	292	146-584	146-584	Begin at the 4-5 leaf stage and then at 10-14 day intervals.
Cabbage	3	438	292	146-584	146-584	Begin at the 4-5 leaf stage and then at 10-14 day intervals.
Cauliflower	3	438	292	146-584	146-584	Begin at the 4-5 leaf stage and then at 10-14 day intervals.

(continued)

Crop	No. of Applications	Broadcast Spray	Band Spray	In Seed Furrow	Transplant Water	Spray Timing
		MI/H	MI/H	MI/H	MI/H	
Corn, Sweet	1	438	365	146-584		2-5 leaf stage.
	2 or more	292	219			Begin at 2-6 leaf stage and then at 7-21 day intervals through the end of tasseling.
Cucumbers	1	584	292			Begin flower bud initiation and first bloom.
	2-3	292	219	146-584	146-584	Begin at flower bud initiation and then 7-10 day intervals.
	3-4	292	219			Begin at transplant, or at the 3-4 leaf stage for direct seeded, and then 7-10 day intervals.
Lettuce	3	438	292	146-584	146-584	Begin at the 4-5 leaf stage and then at 10-14 day intervals.
Melons	1	584	292			Between flower bud initiation and first bloom.
	2-3	292	219	146-584	146-584	Begin at flower bud initiation and then at 7-10 day intervals.
	4-6	219	146			Begin 2 weeks after emergence and then at 7-14 day intervals.
Onions	3	438	292	146-584		Begin 2 weeks after emergence and then at 10-14 day intervals.
Peppers	4-6	292	219	146-584	146-584	Begin at transplant, or at the 3-4 leaf stage for direct seeded, and then at 7-14 day intervals. Do not apply within 14 days of harvest.
Potatoes	1	1168	584	146-584		Tuber initiation.
	3	584	292			Begin at stolon formation (8-10 leaf stage) and then at 10-14 day intervals

(continued)

Crop	No. of Applications	Broadcast Spray	Band Spray	In Seed Furrow	Transplant Water	Spray Timing
		MI/H	MI/H	MI/H	MI/H	
Squash	1	584	292	146-584		Between flower bud initiation and first bloom. Do not apply within 21 days of harvest.
	2-3	292	219			Begin at flower bud initiation and then at 7-10 day intervals. Do not apply within 21 days of harvest.
	4-6	219	146			Begin 2 weeks after emergence and then at 7-14 day intervals. Do not apply within 21 days of harvest.
Tomatoes	1	584	292			Between flower bud initiation and first bloom. Do not apply within 21 days of harvest.
	2-3	292	219	146-584	146-584	Begin at flower bud initiation and then at 7-10 day intervals.
	4-6	219	146			Begin 2 weeks after emergence, or at transplant, and then at 7-14 day intervals

FIELD CROPS

Crop	No. of Applications	Broadcast Spray	Band Spray	In Seed Furrow	Transplant Water	Spray Timing
		MI/H	MI/H	MI/H	MI/H	
Beets, Sugar	1	1168	584	146-584		6-8 leaf stage.
	2-3	584	292			Begin at the 2 leaf stage and then at 7-14 day intervals.
Canola	3	438	292	146-584		Begin at the 3-5 leaf stage and then at 10-14 day intervals.
Corn	1	584	365	146-584		2-6 leaf stage.
Cotton	1	584	292	146-584		Between flower initiation and final bloom
	2-3	292	219			Early bloom and 7-14 days later.
	3-4	292	219			Begin at pinhead square and then at 7-10 day intervals.
Peanuts	4-6	292	219	146-584		Begin 30 days after planting and then at 7-14 day intervals.
Rice	1	584				2-5 leaf stage or panicle initiation
	2	292				2-5 leaf stage or panicle initiation
Sorghum	1	584	365	146-584		2-6 leaf stage.
Soybeans	1	584	365	146-584		3-7 trifoliolate leaf stage (V4-V8)
	2	292	219			3-7 trifoliolate leaf stage (V4-V8) and 10-17 days later.

Crop	No. of Applications	Broadcast Spray	Band Spray	In Seed Furrow	Transplant Water	Spray Timing
		MI/H	MI/H	MI/H	MI/H	
Tobacco	2	584	365		146-584	1 st 40 days after planting 2 nd after topping
Wheat, Barley Oats, Rye	1-2	584	438	146-584		Start at tillering in the fall and/or spring and when 2 to 3 leaves form on main stem.

SMALL FRUITS, VINES AND TREE FRUITS

Bananas:* Apply 300 to 600 ml/hectare in a band around the root mat and repeat every 10 to 14 days for a total of 10 applications.

Pineapple:* Spray after transplanting with 120 to 240 ml/300 liters and repeat every 10 to 14 days or apply in an irrigation system to provide 324 to 648 ml/hectare.

Strawberries:* Begin sprays at first bloom at 292 ml/hectare or broadcast 146 ml/hectare as band sprays directed at the row. Repeat sprays at 2 to 4 week intervals for a total of 3 to 6 sprays. Do not apply within 7 days of harvest.

Oranges:* Apply at 126 ml to 252 ml/100 liters at first bloom and repeat at each flush of new growth.

YOUNG TREES AND ORNAMENTALS

Shrubs, Established:* For increased vigor, rapid growth and healthy plant appearance, spray foliage with transplant solution (2.96 ml Fortified Stimulate Yield Enhancer/7.57 liters of water) to point of run-off two to three times per year.

Flowering Plants (Roses, Azaleas, etc.):* 3-4 year old—using stock solution (2.96 ml Fortified Stimulate Yield Enhancer/7.57 liters of water)—Take 11.83 ml stock solution in 3.79 liters of water and water in around root zone one time per year, preferably early spring.

Sod:*^A To improve growth, heavy rooting, broadcast 0.58 liter/hectare.

1. To speed up regrowth after harvest, broadcast 29.2 ml/hectare.
2. To boost with another 29.2 ml/hectare, broadcast six weeks later.

Turf:*^A For quick "tie down" after laying and to get turf off to a quick start, use Fortified Stimulate Yield Enhancer as follows:

1. Broadcast 5.9 ml/464.5 sq./m and water in.
2. Second application-30 days later-5.9 ml/464.5 sq./m and water in.

^ADo not apply this product through any type of irrigation system.

NOTICE: FORTIFIED STIMULATE YIELD ENHANCER IS NOT A FERTILIZER. USE IN COMBINATION WITH A GOOD FERTILIZER PROGRAM WHERE INDICATED.

GOLF COURSES

Greens: Make an initial treatment with Fortified Stimulate Yield Enhancer to promote root development and protect against "winter kill" using 5.9 ml/464.5-650.32 sq./m². Thereafter, use 29.57 ml/464.5-650.32 sq./m² every 30 days.

Tees: Use Fortified Stimulate Yield Enhancer at 1.48 ml/111.48-139.35 m² area every 30 days to maintain a healthy mass root growth.

Fairways: To establish the necessary root growth to fully utilize applied fertilizer, use Fortified Stimulate Yield Enhancer at 57.14 ml/hectare two times the first year. Thereafter, use 57.14 ml/hectare one to two times a year.

TRANSPLANTS*

STIMULATE YIELD ENHANCER MAY BE USED WHEN TRANSPLANTING ORNAMENTALS AND YOUNG TREES FOR A FAST AND HEALTHY START. Use 2.96 ml/7.57 liters of water stock solution as follows: At time of transplanting

1. Bare (naked) roots - dip or spray with stock solution (2.96 ml Stimulate Yield Enhancer/7.57 liters of water).
2. Balled plants - spray ball at time of transplant.
3. Mist (not run-off) foliage lightly at time of transplant.
4. Using stock solution, apply 9.35 liters Fortified Stimulate Yield Enhancer in furrow/hectare.

SEED TREATMENT*

Use only on seeds for crops listed elsewhere on the label. Do not use treated seed for food, feed or oil purposes. If this product is intended for commercial seed treatment, treated seed must be labeled in accordance with the requirements. If this product is intended for at planting* use, treat only those seeds needed for immediate use and planting. Do not store excess treated seed beyond planting time. Dispose of excess treated seed by burial away from streams and bodies of water. A dye is not required.

For crops listed: Apply 0.75 to 1.5 ml per kg. of seed. Dilute with water and mist spray on seed while mixing. Do not store seed wet as germination can be reduced if not planted soon after treatment.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Product Storage: Do not store in direct sunlight. Avoid freezing temperatures.

Product Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Handling: Use label language appropriate for container size and type.

Non-refillable containers. Do not reuse or refill this container. Clean container promptly after emptying.

Non-refillable container equal to or less than 20 liters. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 120 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by province and local authorities, by burning. If burned, stay out of smoke.

Non-refillable container greater than 20 liters. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill, or by incineration, or allowed by province and local authorities, by burning. If burned, stay out of smoke.

FOR CHEMICAL EMERGENCY:

**Spill, leak, fire, exposure or accident, call
CHEMTREC 1-800-424-9300**

WARRANTY STATEMENT: To the extent consistent with applicable law, Stoller Enterprises, Inc. warrants that the product conforms to the description on the label and is reasonably fit for the purposes set forth on the label, when used according to directions under normal use conditions. Neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use of this product contrary to the label instructions; the buyer assumes the risk of any such uses.

NOTES

Fortified Stimulate Yield Enhancer

ACTIVE INGREDIENTS:

Cytokinin (as kinetin)	0.009%
Gibberellic acid	0.005%
Indole-3-butyric acid	0.005%
Indole-3-acetic acid	0.005%

This product contains: 0.002682 grams cytokinin, 0.001490 grams gibberellic acid, 0.001490 grams indole-3-butyric acid, 0.001490 grams indole-3-acetic acid per fluid ounce.

CONTAINS NON-PLANT FOOD INGREDIENTS:

0.009% Cytokinin
0.005% Gibberellic Acid
0.005% Indole-3-butyric Acid
0.005% Indole-3-acetic Acid

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>

KEEP OUT OF REACH OF CHILDREN
CAUTION

SEE ADDITIONAL PRECAUTIONARY STATEMENTS ON LEFT PANEL.

Registration Number 2018066A Fertilizers Act

Read attached label before using.

FIRST AID

If inhaled	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or ambulance; then give artificial respiration, preferably mouth-to-mouth, if possible.• Call a poison control center or doctor for further treatment advice.
If in eyes	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes; then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
If on skin or clothing	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency information concerning this product, call the National Pesticides Information Center (NPIC) at 1-800-858-7378 seven days a week, 6:30 am to 4:30 pm Pacific Time (NPIC website: www.npic.orst.edu).

Manufactured by

Stoller Enterprises, Inc.

9090 Katy Freeway, Suite 400
Houston, Texas 77024 USA

Phone: 800-539-5283
or 713-461-1493